

The time evolution of the Bohmian Pilot Wave

M. Kozłowski, J. Marciak-Kozłowska

Institute of Electron Technology Al. Lotników 32/46, 02-668 Warsaw,
Poland

Abstract

In this paper the Newton-Schrödinger-Bohm equation is solved for particles with $m > M_P$. It is shown that the Bohmian pilot wave for particles with $m > M_P$ oscillates with frequency $\omega = \tau_P^{-1}$, where τ_P is the Planck time.

Key words: Macroscopic particles; pilot wave.

1 The Newton-Schrödinger-Bohm equation

D. Bohm presented the pilot wave theory in 1952 and de Broglie had presented a similar theory in the mid 1920's. It was rejected in 1950's and the rejection had nothing to do with de Broglie and Bohm later works.

There is always the possibility that the pilot wave has a primitive, mind like property. That's how Bohm described it. We can say that all the particles in the Universe and even Universe have their own pilot waves, their own information. Then the consciousness for example is the very complicated receiver of the surrounding pilot wave fields.

In our paper, <http://lanl.arxiv.org/quant-ph/0402069/> a study of the Newton-Schrödinger-Bohm (NSB) equation for the pilot wave was developed:

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi + V\Psi - \frac{\hbar^2}{2M_P} \nabla^2 \Psi + \frac{\hbar^2}{2M_P} \left(\nabla^2 \Psi - \frac{1}{c^2} \frac{\partial^2 \Psi}{\partial t^2} \right). \quad (1)$$

In Eq. (1) m is the mass of the quantum particle and M_P is the Planck mass ($M_P \approx 10^{-5}$ g).

For elementary particles with mass $m \ll M_P$ we obtain from Eq. (1)

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi + V\Psi + \frac{\hbar^2}{2M_P} \left(\nabla^2 \Psi - \frac{1}{c^2} \frac{\partial^2 \Psi}{\partial t^2} \right) \quad (2)$$

and for macroscopic particles with $m \gg M_P$ equation (1) has the form:

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2M_P} \nabla^2 \Psi + \frac{\hbar^2}{2M_P} \left(\nabla^2 \Psi - \frac{1}{c^2} \frac{\partial^2 \Psi}{\partial t^2} \right) + V\Psi \quad (3)$$

or

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2M_P c^2} \frac{\partial^2 \Psi}{\partial t^2} + V\Psi$$

and is dependent of m .

2 Discussion of the results

In the following we will discuss the pilot wave time evolution for the macroscopic particles, i.e. for particles with $m \gg M_P$.

For $V = \text{const}$ we seek the solution of Eq. (3) in the form

$$\Psi = e^{\gamma t}. \quad (4)$$

After substitution formula (4) to Eq. (3) one's obtains

$$M_P \gamma^2 + \frac{2M_P^2 c^2}{\hbar} \gamma - \frac{2M_P^2 c^2}{\hbar^2} V = 0 \quad (5)$$

with the solution

$$\gamma_{1,2} = -\frac{iM_P c^2}{\hbar} \pm \frac{M_P c^2}{\hbar} \sqrt{-1 + \frac{2V}{M_P c^2}}. \quad (6)$$

For a free particle, $V = 0$ we obtain:

$$\gamma_{1,2} = \begin{cases} 0, \\ -\frac{2M_P c^2}{\hbar} i. \end{cases} \quad (7)$$

According to formulae (4) and (7) equation (3) has the solution

$$\Psi(t) = A + B e^{-\frac{2M_P c^2 i}{\hbar} t}. \quad (8)$$

For $t = 0$ we put $\Psi(0) = 0$, then

$$\Psi(t) = A \left(1 - e^{-\frac{2ti}{\tau_P}} \right),$$

where $\tau_P = \text{Planck time}$

$$\tau = \frac{\hbar}{M_P c^2}. \quad (9)$$

In Fig. 1 $\Re \Psi(t)$ is presented for $t \in (0, 100\tau)$ and $t \in (0, 1000\tau)$ respectively. From Fig. 1 one's can conclude that the free particle in reality is jittering with frequency $\omega = \tau^{-1}$ and quantum energy $E = \hbar\omega$,

$$E = \hbar\omega = 10^{19} \text{GeV} \quad (10)$$

and period T

$$T = 10^{-43} \text{s}.$$

Considering that the contemporary limiting time resolution is of the order 10^{-18} s (attosecond laser pulses) the jittering of the free macroscopic particle can not be observed.

3 Conclusion

The NSB equation offers the new insight into the realm of the subquantum structure of the space-time. It occurs that free macroscopic particles, with mass $m > M_P$ are embedded in the field of the oscillations with the period of $T \approx 10^{-43}$ s and energy $E \approx 10^{19}$ GeV.





Figure 1: